

“Наиболее удобной моделью для отработки
такой связи является цепочка Тоды”

Л.Д. Фаддеев, Препринт ЛОМИ Р-2-79

Quantum Toda Chains Intertwined

Anton Gerasimov, Dimitri Lebedev, Sergey Oblezin

To Ludvig Dmitrievich Faddeev on the occasion of his 75-th birthday

Abstract

Abstract. We conjecture an explicit construction of integral operators intertwining various quantum Toda chains. Compositions of the intertwining operators provide recursive and $\mathcal{Q}$ -operators for quantum Toda chains. In particular we propose a generalization of our previous results for Toda chains corresponding to classical Lie algebra to the generic BC_n and Inozemtsev Toda chains. We also conjecture explicit form of $\mathcal{Q}$ -operators for closed Toda chains corresponding to Lie algebras B_∞ , C_∞ , D_∞ , affine Lie algebras $B_n^{(1)}$, $C_n^{(1)}$, $D_n^{(1)}$, $D_n^{(2)}$, $A_{2n-1}^{(2)}$, $A_{2n}^{(2)}$ and the affine analogs of BC_n and Inozemtsev Toda chains.

Introduction

An interesting integral representation for common eigenfunctions of the quantum A_n -Toda chain Hamiltonian operators was proposed in [Gi]. Using the representation theory approach to Toda chain eigenfunctions the representation [Gi] was generalized to other classical series B_n , C_n and D_n [GLO2]. In [GKLO], [GLO2] we stress an important role of the elementary intertwining operators in the construction of solutions of quantum Toda chains. The common eigenfunctions of quantum Toda Hamiltonians corresponding to the same classical series can be obtained by a recursive application of some integral operators. In general the recursive integral operators can be naturally represented as a product of elementary integral operators intertwining quantum Toda Hamiltonians corresponding to different classical series. The integral kernels of the elementary intertwining operators have a uniform structure of the exponent of a linear combination of exponents in natural coordinates. This decomposition of the recursive operators into elementary intertwiners seems has a methodological importance for quantization of classical Toda chains. Indeed, for classical Toda chains the recursive operators are given by complicated canonical transformations. The direct quantization of these canonical transformations is far from obvious. Let classical canonical transformations be naturally split into products of elementary canonical transformations allowing standard quantization. Then the quantization of the canonical transformation is easily obtained as a product of quantized elementary canonical transformations. As it was argued in [GLO2] and is demonstrated in this paper this is precisely the case for Toda chains.

One can expect that the algebra of intertwining operators acting between various Toda chains should have an interesting interpretation in representation theory of classical series of Lie groups. The representation theory interpretation of a particular instance of the integral intertwining operator commuting with quantum Toda chain Hamiltonians was proposed in [GLO3]. These operators were identified with the generators of Hecke algebras of the classical Lie groups with respect to maximal compact subgroups. The affine version of these operators were introduced long ago by Pasquier and Gaudin as $\mathcal{Q}$ -operator for affine Toda chain [PG].

In this paper we extend the integral intertwiner operator approach to open Toda chains whose representation theory interpretation for generic coupling constants is not known. These include BC_n -Toda chains constructed by Sklyanin [S] and Toda chains proposed by Inozemtsev in the classical setting [I]. In the following we refer to the latter series of Toda chains as I_n -Toda chains. We propose elementary intertwining operators for BC_n and I_n Toda chains and check the intertwining properties with respect to quadratic quantum Hamiltonians. The elementary intertwiners appear to relate BC and I -Toda chains thus stressing the importance of the Inozemtsev Toda chain series as an honorable member of the Toda chains family. The constructed elementary intertwiners allow us to propose explicit integral expressions for common eigenfunctions of BC_n - and I_n -Toda chain Hamiltonians for generic spectrum.

In the second part of the paper we propose analogs of the Pasquier-Gaudin integral $\mathcal{Q}$ -operator [PG] for affine Toda chains based on classical series of affine Lie algebras $A_n^{(1)}$, $A_{2n}^{(2)}$, $A_{2n-1}^{(2)}$, $B_n^{(1)}$, $C_n^{(1)}$, $D_n^{(1)}$, $D_n^{(2)}$ infinite Lie algebras B_∞ , C_∞ and D_∞ , affine and semi-infinite version of BC_n and I_n Toda-chains.

Finally let us stress that the explicit integral expressions for common eigenfunctions constructed using elementary intertwiners might provide a general approach to the quantization of classical integrable systems. This program was completely realized for A_n -Toda chains in [GLO3]. The first step is a construction of recursive operators as a product of quantized elementary intertwining integral operators and checking commutation relations with quadratic quantum Hamiltonian. The next step is a construction of quadratic Hamiltonian eigenfunctions using the recursive operators. Finally one should check orthogonality and completeness relations for the constructed eigenfunctions. Note that the complete set of eigenfunctions can be considered (after a standard Fourier transform) as an integral kernel of a unitary transformation U intertwining quantum Toda chain with the quantum theory of free particles (i.e. having zero potential). The higher quantum Hamiltonians can now be obtained by a conjugation by U of the standard set of the higher quantum Hamiltonians for a collection of free particles. The correct classical limit of these Hamiltonians is guaranteed by the correct classical limit of the quantum recursive operators used to construct U . It would be interesting to apply this line of reasoning to BC_n - and I_n -Toda chains which do not allow simple representation theory realization. What is missed at the moment are the orthogonality and completeness relations for these sets of functions. The proof of the orthogonality and completeness relations consists of routine manipulations with iterative integrals and will be published elsewhere. Modern approach to solving quantum integrable systems based on the Quantum Inverse Scattering Method (QISM) achieved a spectacular success for various classes of quantum integrable models (see [F] for a concise account). One might hope that the approach based on the explicit integral operators used in this paper might add some complimentary information about the structure of quantum integrable systems.

This paper is an updated version of the earlier note [GLO1]. After [GLO1], [GLO2] the paper [KS] with the results on classical recursion operators for BC_n -Toda chains appeared. We expect that the classical limit of our constructions for BC_n Toda chain should be compatible with the

results of [KS].

The plan of this paper is as follows. In Section 1 we summarize the results of [Gi], [GKLO], [GLO2] for the case of zero spectrum. In Section 2 we propose elementary integral intertwining operators and integral representations for BC_n - and I_n -Toda chain eigenfunctions. In Section 3 analogs of Pasquier-Gaudin Q -operator at zero spectral parameter are proposed for affine versions of Toda chains corresponding to classical Lie groups and BC_n - and I_n -Toda chains. In Section 4 the analogous conjectures are stated for semi-infinite Toda chains.

Acknowledgments: The research was supported by grants RFBR-08-01-00931-a, 09-01-93108-HIHHIЛ-a, AG was also partly supported by Science Foundation Ireland grant.

1 Integral representation for the eigenfunctions of quantum Toda chains for classical groups

We start reviewing a construction of the intertwining operators and common eigenfunction of quantum Toda chains corresponding to classical Lie algebras [GLO2]. Then we conjecture explicit integral representations for eigenfunctions for generic BC_n and I_n Toda chains.

Let us first recall the standard facts on Toda chains corresponding to arbitrary root systems (see e.g. [RSTS]). Let $\mathfrak{g}$ be a simple Lie algebra, $\mathfrak{h}$ be a Cartan subalgebra, $n = \dim \mathfrak{h}$ be the rank of $\mathfrak{g}$, $R \subset \mathfrak{h}^*$ be the root system, W be the Weyl group. Let us fix a decomposition $R = R_+ \cup R_-$ of the roots on positive and negative roots. Let $\alpha_1, \dots, \alpha_n$ be the bases of simple roots. Let $(,)$ be a W -invariant bilinear symmetric form on $\mathfrak{h}^*$ normalized so that $(\alpha, \alpha) = 2$ for a long root. This form provides an identification of $\mathfrak{h}$ with $\mathfrak{h}^*$ and thus can be considered as a bilinear form on $\mathfrak{h}$. Choose an orthonormal basis $e = \{e_1, \dots, e_n\}$ in $\mathfrak{h}$. Then for any $x \in \mathfrak{h}$ one has a decomposition $x = \sum_{i=1}^n x_i e_i$. One associates with these data an open Toda chain with a quadratic Hamiltonian

$$H_2^R(\underline{x}) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} + \sum_{i=1}^n g_i e^{\alpha_i(x)} \quad (1.1)$$

where g_i are (in general complex) coupling constants. This Hamiltonian is a one of the generators $\mathcal{H}_i^R$, $i = 1, 2, \dots, \text{rank}(R)$ of the commutative ring of quantum Hamiltonians of the corresponding Toda chain. The common eigenfunctions of the quantum Hamiltonians satisfy the equations

$$H_i^R(\underline{x}) \Psi_{\underline{\lambda}}^R(\underline{x}) = c_i(\underline{\lambda}) \Psi_{\underline{\lambda}}^R(\underline{x}),$$

where $c_i(\underline{\lambda})$ are polynomials of the complex variables $\underline{\lambda} = (\lambda_1, \dots, \lambda_n)$ parameterizing spectrum. In the following we will consider only the eigenfunctions corresponding to zero eigenvalues of the Hamiltonians and omit the index $\underline{\lambda}$. Borrowing Physics terminology we will call such eigenfunctions ground state wave functions.

1.1 Integral representation of A_{n-1} -Toda chain eigenfunctions

In this Subsection we recall a recursive construction [GKLO] of the Givental integral representation of $\mathfrak{gl}_n$ -Toda chain eigenfunctions [Gi]. Quantum $\mathfrak{gl}_n$ -Toda chain is characterized by its linear and

quadratic Hamiltonians

$$\begin{aligned} H_1^{\mathfrak{gl}_n}(\underline{x}) &= \imath \sum_{i=1}^n \frac{\partial}{\partial x_i}, \\ H_2^{\mathfrak{gl}_n}(\underline{x}) &= -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} + \sum_{i=1}^{n-1} g_i e^{x_{i+1}-x_i}, \end{aligned} \quad (1.2)$$

where $\underline{x} = (x_1, \dots, x_n)$ and g_i are coupling constants. These Hamiltonians are the first two of a series of generators $H_i^{\mathfrak{gl}_n}$, $i = 1, 2, \dots, n$ of the commutative ring of quantum Hamiltonians of $\mathfrak{gl}_n$ -Toda chain. The common eigenfunctions of the quantum Hamiltonians satisfy the equations

$$H_i^{\mathfrak{gl}_n}(\underline{x}) \Psi_{\underline{\lambda}}^{\mathfrak{gl}_n}(\underline{x}) = c_i(\underline{\lambda}) \Psi_{\underline{\lambda}}^{\mathfrak{gl}_n}(\underline{x}), \quad (1.3)$$

where $c_i(\underline{\lambda})$ are the symmetric polynomials of the complex variables $\underline{\lambda} = (\lambda_1, \dots, \lambda_n)$ parameterizing spectrum. For instance $c_1(\underline{\lambda}) = \lambda_1 + \dots + \lambda_n$ and $c_2(\underline{\lambda}) = \lambda_1^2 + \dots + \lambda_n^2$. Common eigenfunctions for $\mathfrak{sl}_n$ -Toda chain are obtained by imposing the additional condition $c_1(\underline{\lambda}) = 0$ in (1.3). In the following we will consider ground state wave functions corresponding to $\underline{\lambda} = 0$ and thus will not distinguish $\mathfrak{gl}_n$ - and $\mathfrak{sl}_n$ -Toda chains.

1.1.1 The following integral representation for a ground state wave function quantum Hamiltonians of $\mathfrak{gl}_n$ -Toda chain was derived in [Gi]

$$\Psi^{\mathfrak{gl}_n}(\underline{x}) = \int_{\Gamma} e^{\mathcal{F}_n(x)} \prod_{k=1}^{n-1} \prod_{i=1}^k dx_{k,i}, \quad (1.4)$$

where $x_{n,i} := x_i$, $\underline{x} = (x_1, \dots, x_n)$, the function $\mathcal{F}_n(x)$ is given by

$$\mathcal{F}_n(x) = - \sum_{k=1}^{n-1} \sum_{i=1}^k \left(e^{x_{k,i}-x_{k+1,i}} + g_i e^{x_{k+1,i+1}-x_{k,i}} \right), \quad (1.5)$$

and the integration domain Γ is a middle dimensional submanifold in the $n(n-1)/2$ - dimensional complex torus with coordinates $\{\exp x_{k,i}, i \leq 1 \leq k \leq n-1\}$. The choice of the integration contour is dictated by the condition on the integrand to decrease exponentially at infinities.

The integral representation (1.4) can be derived using the recursion over the rank n of the Lie algebra $\mathfrak{gl}_n$ [GKLO]. Indeed (1.4) is naturally represented in the recursive form

$$\Psi^{\mathfrak{gl}_n}(\underline{x}) = \int_{\Gamma} \prod_{k=1}^{n-1} Q_{k+1,k}(\underline{x}_{k+1}; \underline{x}_k) d\underline{x}_k, \quad (1.6)$$

with the integral kernel

$$Q_{k+1,k}(\underline{x}_{k+1}; \underline{x}_k) = \exp \left\{ - \sum_{i=1}^k \left(e^{x_{k,i}-x_{k+1,i}} + g_i e^{x_{k+1,i+1}-x_{k,i}} \right) \right\}. \quad (1.7)$$

Here we define $\underline{x}_k = (x_{k,1}, \dots, x_{k,k})$ and $x_i := x_{n,i}$, $1 \leq i \leq n$. The zero eigenvalue property of (1.4) with respect to quantum Hamiltonians is then translated into the intertwining relations between integral operators with the kernels $Q_{k+1,k}$ and Hamiltonians $H_i^{\mathfrak{gl}_k}$ and $H_i^{\mathfrak{gl}_{k+1}}$. In particular for the quadratic Hamiltonian (1.2) we have the following easily verified property

$$H^{\mathfrak{gl}_{k+1}}(\underline{x}_{k+1}) Q_{k+1,k}(\underline{x}_{k+1}, \underline{x}_k) = Q_{k+1,k}(\underline{x}_{k+1}, \underline{x}_k) H^{\mathfrak{gl}_k}(\underline{x}_k), \quad (1.8)$$

where

$$H^{\mathfrak{gl}_k}(\underline{x}_k) = -\frac{1}{2} \sum_{i=1}^k \frac{\partial^2}{\partial x_{k,i}^2} + \sum_{i=1}^{k-1} g_i e^{x_{k,i+1}-x_{k,i}}. \quad (1.9)$$

Here and in the following we assume that in the relations similar to (1.8) the Hamiltonian operator on l.h.s. acts on the right and the Hamiltonian on r.h.s. acts on the left. From (1.8) one can easily deduce

$$\left(-\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} + \sum_{i=1}^{n-1} g_i e^{x_{i+1}-x_i} \right) \Psi^{\mathfrak{gl}_n}(\underline{x}) = 0. \quad (1.10)$$

The eigenvalue property for higher Hamiltonians is not so easy to verify and deserve more involved considerations [Gi], [GKLO].

1.1.2 The integral operator defined by the kernel (1.7) is closely related with a $\mathcal{Q}$ -operator realizing Bäcklund transformations in a Toda chain corresponding to affine Lie algebra $\widehat{\mathfrak{gl}}_n$. This is a particular example of the integral intertwining operator introduced by Pasquier and Gaudin [PG]. The integral kernel is given by

$$Q^{\widehat{\mathfrak{gl}}_n}(\underline{x}, \underline{y}) = \exp \left\{ - \sum_{i=1}^n \left(e^{x_i-y_i} + g_i e^{y_{i+1}-x_i} \right) \right\}, \quad (1.11)$$

where $x_{i+n} = x_i$ and $y_{i+n} = y_i$. This operator commutes with the Hamiltonian operators of the closed Toda chain. Thus for example for the quadratic Hamiltonian we have

$$H_2^{\widehat{\mathfrak{gl}}_n}(\underline{x}) Q^{\widehat{\mathfrak{gl}}_n}(\underline{x}, \underline{y}) = Q^{\widehat{\mathfrak{gl}}_n}(\underline{x}, \underline{y}) H_2^{\widehat{\mathfrak{gl}}_n}(\underline{y}), \quad (1.12)$$

where

$$H_2^{\widehat{\mathfrak{gl}}_n}(\underline{x}) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} + \sum_{i=1}^n g_i e^{x_{i+1}-x_i}. \quad (1.13)$$

Here we impose the conditions $x_{i+n} = x_i$. The recursive operator (1.7) can be obtained from the operator (1.11) in the limit $g_n \rightarrow 0$, $x_n \rightarrow -\infty$ in such a way that $g_n e^{-x_n} \rightarrow 0$.

1.2 Toda chains corresponding to classical Lie algebras

In this Subsection we recall the explicit form of the quadratic quantum Hamiltonians of Toda chains corresponding to classical Lie algebras.

1.2.1 Let $e = \{e_1, \dots, e_n\}$ be an orthonormal basis in $\mathbb{R}^n$. The root system of type B_n can be defined as

$$\alpha_1 = e_1, \quad \alpha_{i+1} = e_{i+1} - e_i, \quad 1 \leq i \leq n-1, \quad (1.14)$$

and the corresponding Dynkin diagram is

$$\alpha_1 \Longleftarrow \alpha_2 \text{ --- } \dots \text{ --- } \alpha_{n-1} \text{ --- } \alpha_n. \quad (1.15)$$

Quadratic Hamiltonian operator of the corresponding Toda chain is given by

$$H_2^{B_n}(\underline{x}) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} + g_1 e^{x_1} + \sum_{i=1}^{n-1} g_{i+1} e^{x_{i+1} - x_i}. \quad (1.16)$$

1.2.2 The root system of type C_n can be defined as

$$\alpha_1 = 2e_1, \quad \alpha_{i+1} = e_{i+1} - e_i, \quad 1 \leq i \leq n-1, \quad (1.17)$$

and the corresponding Dynkin diagram is

$$\alpha_1 \implies \alpha_2 \text{ --- } \cdots \text{ --- } \alpha_{n-1} \text{ --- } \alpha_n. \quad (1.18)$$

Quadratic Hamiltonian operator of the corresponding Toda chain is given by

$$H^{C_n}(\underline{z}) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial z_i^2} + 2g_1 e^{2z_1} + \sum_{i=1}^{n-1} g_{i+1} e^{z_{i+1} - z_i}, \quad (1.19)$$

where $\underline{z} = (x_1, \dots, z_n)$.

There is a more general Toda chain associated with root system C_n (see e.g. [RSTS], Remark p.61) with the quadratic Hamiltonian


$$H_2^{BC_n^*}(\underline{z}) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial z_i^2} - \frac{g_1}{2} e^{z_1} + \frac{g_1^2}{2} e^{2z_1} + \sum_{i=1}^{n-1} g_{i+1} e^{z_{i+1} - z_i}, \quad (1.20)$$

The BC_n^* -Toda chain can be obtained as a degeneration of the generic BC_n -Toda chain discussed in the next. This explains our notations.

1.2.3 The root system of type D_n is

$$\alpha_1 = e_1 + e_2, \quad \alpha_{i+1} = e_{i+1} - e_i, \quad 1 \leq i \leq n-1, \quad (1.21)$$

and the corresponding Dynkin diagram is



Quadratic Hamiltonian operator of the D_n -Toda chain is given by

$$H^{D_n}(\underline{x}) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} + g_1 g_2 e^{x_1 + x_2} + \sum_{i=1}^{n-1} g_{i+1} e^{x_{i+1} - x_i}. \quad (1.22)$$

1.3 Intertwining integral operators

We start with the elementary intertwining operators relating Toda chains corresponding to different series. These elementary intertwiners have simple integral kernels given by exponents of the linear combination of exponents. The integration contours are chosen in such way that the integrals considered below converge.

1.3.1 Integral kernel

$$Q_{B_{n-1}}^{BC_n^*}(\underline{z}_n; \underline{x}_{n-1}) = \exp \left\{ - \left(g_1 e^{z_{n,1}} + \sum_{i=1}^{n-1} \left(e^{x_{n-1,i} - z_{n,i}} + g_{i+1} e^{z_{n,i+1} - x_{n-1,i}} \right) \right) \right\}, \quad (1.23)$$

satisfies the following relations

$$H^{BC_n^*}(\underline{z}_n) Q_{B_{n-1}}^{BC_n^*}(\underline{z}_n, \underline{x}_{n-1}) = Q_{B_{n-1}}^{BC_n^*}(\underline{z}_n, \underline{x}_{n-1}) H^{B_{n-1}}(\underline{x}_{n-1}). \quad (1.24)$$

Thus the corresponding integral operator intertwines quadratic Hamiltonians of BC_n^* - and B_{n-1} -Toda chains. Similarly an elementary operator intertwining quadratic B_n and BC_n^* Hamiltonians has an integral kernel

$$Q_{BC_n^*}^{B_n}(\underline{x}_n; \underline{z}_n) = \exp \left\{ - \left(g_1 e^{z_{n,1}} + \sum_{i=1}^{n-1} \left(e^{x_{n,i} - z_{n,i}} + g_{i+1} e^{z_{n,i+1} - x_{n,i}} \right) + e^{x_{nn} - z_{nn}} \right) \right\}. \quad (1.25)$$

1.3.2 An integral operator intertwining C_n and D_n Hamiltonians has the kernel

$$Q_{D_n}^{C_n}(\underline{z}_n; \underline{x}_n) = \exp \left\{ - \left(g_1 e^{x_{n,1} + z_{n,1}} + \sum_{i=1}^{n-1} \left(e^{z_{n,i} - x_{n,i}} + g_{i+1} e^{x_{n,i+1} - z_{n,i}} \right) + e^{z_{nn} - x_{nn}} \right) \right\}. \quad (1.26)$$

Similarly an integral operator with the kernel

$$Q_{C_{n-1}}^{D_n}(\underline{x}_n; \underline{z}_{n-1}) = \exp \left\{ - \left(g_1 e^{x_{n,1} + z_{n-1,1}} + \sum_{i=1}^{n-1} e^{z_{n-1,i} - x_{n,i}} + g_{i+1} e^{x_{n,i+1} - z_{n-1,i}} \right) \right\}, \quad (1.27)$$

intertwines the following D_n and C_{n-1} quadratic Hamiltonians

$$H^{C_{n-1}}(\underline{z}_{n-1}) \cdot Q_{C_{n-1}}^{D_n}(\underline{x}_n; \underline{z}_{n-1}) = Q_{C_{n-1}}^{D_n}(\underline{x}_n; \underline{z}_{n-1}) H^{D_n}(\underline{x}_n). \quad (1.28)$$

To describe a recursion structure of the ground state eigenfunction of C_n -Toda chain let us introduce the integral operators $Q_{C_{k-1}}^{C_k}$ with the kernels given for $k = 2, \dots, n$ by the convolutions of the intertwining operators $Q_{C_k}^{D_k}$ and $Q_{D_k}^{C_{k-1}}$:

$$Q_{C_{k-1}}^{C_k}(\underline{z}_k; \underline{z}_{k-1}) = \int_{\Gamma^{(k)}} Q_{D_k}^{C_k}(\underline{z}_k; \underline{x}_k) Q_{C_{k-1}}^{D_k}(\underline{x}_k; \underline{z}_{k-1}) d\underline{x}_k. \quad (1.29)$$

For $k = 1$, we set

$$Q_{C_0}^{C_1}(z_{1,1}) := \Psi^{C_1}(z_{1,1}) = \int_{\Gamma^{(1)}} \exp \{ -g_1 e^{x_{1,1} + z_{1,1}} - e^{z_{1,1} - x_{1,1}} \} dx_{1,1}. \quad (1.30)$$

1.3.3 For $k = 2, \dots, n$ introduce the integral operators $Q_{D_{k-1}}^{D_k}$ with kernels given by the convolutions of the kernels of the intertwining operators $Q_{C_{k-1}}^{D_k}$ and $Q_{D_{k-1}}^{C_{k-1}}$

$$Q_{D_{k-1}}^{D_k}(\underline{x}_k; \underline{x}_{k-1}) = \int_{\Gamma^{(k)}} Q_{C_{k-1}}^{D_k}(\underline{x}_k; \underline{z}_{k-1}) Q_{D_{k-1}}^{C_{k-1}}(\underline{z}_{k-1}; \underline{x}_{k-1}) d\underline{z}_{k-1}. \quad (1.31)$$

For $k = 1$ we set $Q_{D_0}^{D_1}(x_{1,1}) = 1$.

1.3.4 Denote $\underline{x}_k = (x_{k,1}, \dots, x_{k,k})$, $\underline{z}_k = (z_{k,1}, \dots, z_{k,k})$. Let us introduce integral operators $Q_{B_{k-1}}^{B_k}$ for $k = 2, \dots, n$ with the kernels given by the convolutions of the kernels of elementary intertwiners $Q_{BC_k^*}^{B_k}$ and $Q_{B_{k-1}}^{BC_k^*}$:

$$Q_{B_{k-1}}^{B_k}(\underline{x}_k; \underline{x}_{k-1}) = \int_{\Gamma^{(k)}} Q_{BC_k^*}^{B_k}(\underline{x}_k; \underline{z}_k) Q_{B_{k-1}}^{BC_k^*}(\underline{z}_k; \underline{x}_{k-1}) d\underline{z}_k. \quad (1.32)$$

For $k = 1$ we set

$$Q_{B_0}^{B_1}(x_{1,1}) := \Psi^{B_1}(x_{1,1}) = \int_{\Gamma^{(1)}} \exp\{-2g_1 e^{z_{1,1}} - e^{x_{1,1} - z_{1,1}}\} dz_{1,1}.$$

1.4 Integral representations of Toda chain eigenfunctions

Using the integral intertwining operators discussed above we can construct recursively ground state eigenfunctions for quantum Toda chains associated with classical Lie algebras. Thus obtained integral representations are compatible with the representation theory construction of the of generic eigenfunctions for classical Lie algebras [GLO2]. The eigenfunction property with respect to quadratic Hamiltonians follows from the intertwining relations considered in the previous Subsection. For higher quantum Hamiltonians we should at the moment rely on the results obtained in representation theory approach.

Proposition 1.1 *The ground state eigenfunction for B_n -Toda chain has the following integral representation*

$$\Psi^{B_n}(x_1, \dots, x_n) = \int_{\Gamma} \prod_{k=1}^n Q_{B_{k-1}}^{B_k}(\underline{x}_k; \underline{x}_{k-1}) \prod_{k=1}^{n-1} d\underline{x}_k, \quad (1.33)$$

where $x_i := x_{n,i}$ and $d\underline{x}_k = dx_{k,1} \cdots dx_{k,k}$. The eigenfunction is given by the integral over a non-compact domain Γ of the real dimension equal to the complex dimension of the generalized flag manifold $X = G/B$, where $G = SO(2n+1, \mathbb{C})$ and B is a Borel subgroup

$$\sum_{k=1}^n (2k-1) = n^2 = |R_+|. \quad (1.34)$$

The domain of integration should be chosen in such a way that the integrand decreases exponentially at infinities.

Proposition 1.2 *The ground state for C_n -Toda chain has the following integral representation*

$$\Psi^{C_n}(z_1, \dots, z_n) = \int_{\Gamma} \prod_{k=1}^n Q_{C_{k-1}}^{C_k}(\underline{z}_k; \underline{z}_{k-1}) \prod_{k=1}^{n-1} d\underline{z}_k, \quad (1.35)$$

where $z_i := z_{n,i}$ and $d\underline{z}_k = dz_{k1} \cdots dz_{kk}$. The eigenfunction is given by the integral over a non-compact domain Γ of the real dimension equal to complex dimension of the generalized flag manifold $X = G/B$, where $G = SO(2n+1, \mathbb{C})$ and B is a Borel subgroup

$$\sum_{k=1}^n (2k-1) = n^2 = |R_+|. \quad (1.36)$$

The domain of integration should be chosen in such a way that the integrand decreases exponentially at infinities.

Proposition 1.3 *The ground state eigenfunction for D_n -Toda chain has the following integral representation*

$$\Psi^{D_n}(x_1, \dots, x_n) = \int_{\Gamma} \prod_{k=1}^n Q_{D_{k-1}}^{D_k}(\underline{x}_k; \underline{x}_{k-1}) \prod_{r=1}^{n-1} d\underline{x}_r, \quad (1.37)$$

where $x_i := x_{n,i}$. The eigenfunction is given by the integral over a non-compact domain Γ of the real dimension equal to complex dimension of the generalized flag manifold $X = G/B$, where $G = SO(2n, \mathbb{C})$ and B is a Borel subgroup

$$\sum_{k=1}^{n-1} 2k = n(n-1) = |R_+|. \quad (1.38)$$

The domain of integration should be chosen in such a way that the integrand decreases exponentially at infinities.

1.5 Quadratic Hamiltonians of BC_n - and I_n - Toda chains

Here we write down quadratic Hamiltonians of the generic BC_n - and I_n -Toda chains.

1.5.1 Non-reduced root system of type BC_n can be defined as

$$\alpha_0 = 2e_1, \quad \alpha_1 = e_1, \quad \alpha_{i+1} = e_{i+1} - e_i, \quad 1 \leq i \leq n-1 \quad (1.39)$$

and the corresponding Dynkin diagram is

$$\begin{array}{c} \alpha_0 \\ \parallel \\ \parallel \\ \parallel \\ \parallel \\ \parallel \\ \downarrow \\ \alpha_1 \end{array} \begin{array}{c} \searrow \\ \searrow \end{array} \begin{array}{c} \alpha_2 \text{ --- } \cdots \text{ --- } \alpha_{n-1} \text{ --- } \alpha_n \end{array} \quad (1.40)$$

where the two vertexes connected by a double dash line correspond to a reduced $\alpha_1 = e_1$ and non-reduced $\alpha_0 = 2e_1 = 2\alpha_1$ roots.

The corresponding BC_n -Toda chain has the following quadratic Hamiltonian

$$H^{BC_n}(\underline{x}) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} + g_1 e^{x_1} + g_2 e^{2x_1} + \sum_{i=1}^{n-1} g_{i+2} e^{x_{i+1}-x_i}, \quad (1.41)$$

where $\underline{x} = (x_1, \dots, x_n)$. Note that quadratic Hamiltonian (1.20) of the BC_n^* -Toda chain is obtained by the following specialization of the coupling constants $g_1 = -g/2$, $g_2 = g^2/2$.

1.5.2 The quadratic Hamiltonian of the I_n -Toda chain [I] is given by

$$H^{I_n}(\underline{x}) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} + \frac{\tilde{g}_1}{(e^{x_1/2} - e^{-x_1/2})^2} + \frac{\tilde{g}_2}{(e^{x_1} - e^{-x_1})^2} + \tilde{g}_3 e^{x_1+x_2} + \sum_{i=1}^{n-1} \tilde{g}_{i+2} e^{x_{i+1}-x_i}, \quad (1.42)$$

where $\underline{x} = (x_1, \dots, x_n)$ and g_i , $i = 1, \dots, n+1$ are generic coupling constants.

1.5.3 The following specialization of the coupling constants $\tilde{g}_i$, $1 \leq i \leq n+1$ will be important in the following

$$\tilde{g}_1 = -\frac{g_1}{\sqrt{2g_2}}, \quad \tilde{g}_2 = 2\frac{g_1}{\sqrt{2g_2}} \left(\frac{g_1}{\sqrt{2g_2}} + 1 \right), \quad \tilde{g}_3 = g_3 \sqrt{\frac{g_2}{2}},$$

and $\tilde{g}_{i+2} = g_{i+2}$ for $1 < i \leq n-1$. Denote the Hamiltonian with the specialized coupling constants by $H^{I_n^*}$:

$$H^{I_n^*}(\underline{z}_n) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial z_{n,i}^2} - \frac{\frac{g_1}{\sqrt{2g_2}}}{(e^{-z_{n,1}/2} - e^{z_{n,1}/2})^2} + \frac{2\frac{g_1}{\sqrt{2g_2}} \left(\frac{g_1}{\sqrt{2g_2}} + 1 \right)}{(e^{-z_{n,1}} - e^{z_{n,1}})^2} + g_3 \sqrt{\frac{g_2}{2}} \left(e^{z_{n,1}+z_{n,2}} + e^{z_{n,2}-z_{n,1}} \right) + \sum_{i=2}^{n-1} g_{i+2} e^{z_{n,i+1}-z_{n,i}}. \quad (1.43)$$

1.6 Integral intertwining operators

Now we conjecture the BC_n and I_n analogs of the intertwining integral operators. The intertwining relations with quadratic quantum Hamiltonians are checked straightforwardly. Hopefully similar intertwining relations hold for higher quantum Hamiltonians.

Lemma 1.1 *An integral operator defined by the following kernel:*

$$Q_{I_n^*}^{BC_n}(\underline{x}_n, \underline{z}_n) = \left(\frac{1 - e^{z_{n,1}}}{1 + e^{z_{n,1}}} \right)^{\frac{g_1}{\sqrt{2g_2}}} \cdot \exp \left\{ - \left(\sqrt{\frac{g_2}{2}} \left(e^{x_{n,1}+z_{n,1}} + e^{x_{n,1}-z_{n,1}} \right) + \sum_{i=2}^n g_{i+1} e^{z_{n,i}-x_{n,i-1}} + e^{x_{n,i}-z_{n,i}} \right) \right\}, \quad (1.44)$$

intertwines quadratic BC_n and I_n^* Hamiltonians (1.41) and (1.43)

$$H^{BC_n}(\underline{x}_n) \cdot Q_{I_n^*}^{BC_n}(\underline{x}_n, \underline{z}_n) = Q_{I_n^*}^{BC_n}(\underline{x}_n, \underline{z}_n) \cdot H^{I_n^*}(\underline{z}_n). \quad (1.45)$$

Lemma 1.2 Let $Q_{I_{n+1}^*}^{BC_n}$ be an integral operator with the kernel

$$Q_{I_{n+1}^*}^{BC_n}(\underline{x}_n, \underline{z}_{n+1}) = \left(\frac{1 - e^{z_{n+1,1}}}{1 + e^{z_{n+1,1}}} \right)^{\frac{g_1}{\sqrt{2g_2}}} \cdot \exp \left\{ - \left(\sqrt{\frac{g_2}{2}} \left(e^{x_{n,1}+z_{n+1,1}} + e^{x_{n,1}-z_{n+1,1}} \right) + \sum_{i=2}^n \left(g_{i+1} e^{z_{n+1,i}-x_{n,i-1}} + e^{x_{n,i}-z_{n+1,i}} \right) + g_{n+2} e^{z_{n+1,n+1}-x_{nn}} \right) \right\}. \quad (1.46)$$

The following intertwining relation between the quadratic quantum BC_n and I_{n+1}^* Hamiltonians holds

$$H^{BC_n}(\underline{x}_n) \cdot Q_{I_{n+1}^*}^{BC_n}(\underline{x}_n, \underline{z}_{n+1}) = Q_{I_{n+1}^*}^{BC_n}(\underline{x}_n, \underline{z}_{n+1}) \cdot H^{I_{n+1}^*}(\underline{z}_{n+1}). \quad (1.47)$$

Let us denote the kernel of the inverse integral transformation by

$$Q_{BC_n}^{I_{n+1}^*}(\underline{z}_{n+1}, \underline{x}_n) := Q_{I_{n+1}^*}^{BC_n}(\underline{x}_n, \underline{z}_{n+1}).$$

For any $k = 2, \dots, n$ define the integral operators $Q_{BC_k}^{BC_{k-1}}$ as convolutions of the kernels of elementary intertwiners $Q_{I_k^*}^{BC_k}$ and $Q_{BC_{k-1}}^{I_k^*}$:

$$Q_{BC_{k-1}}^{BC_k}(\underline{x}_k; \underline{x}_{k-1}) = \int_{\Gamma^{(k)}} Q_{I_k^*}^{BC_k}(\underline{x}_k; \underline{z}_k) Q_{BC_{k-1}}^{I_k^*}(\underline{z}_k; \underline{x}_{k-1}) d\underline{z}_k. \quad (1.48)$$

For $k = 1$ we set

$$Q_{BC_0}^{BC_1}(x_{1,1}) = \int_{\Gamma^{(1)}} \left(\frac{1 - e^{z_{11}}}{1 + e^{z_{11}}} \right)^{\frac{2g_1}{\sqrt{2g_2}}} \exp\{-\sqrt{g_2/2}(e^{x_{11}+z_{11}} + e^{x_{11}-z_{11}})\} dz_{11}. \quad (1.49)$$

Let us define the following quadratic BC and I Hamiltonians

$$H^{BC_n}(\underline{x}) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} + g_1 e^{x_1} + g_2 e^{2x_1} + \sum_{k=1}^{n-1} g_{k+2} e^{x_{k+1}-x_k}, \quad (1.50)$$

$$H^{I_n}(\underline{z}) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial z_i^2} + \frac{\tilde{g}_1}{(e^{-z_1/2} - e^{z_1/2})^2} + \frac{\tilde{g}_2}{(e^{-z_1} - e^{z_1})^2} + g_3 \sqrt{\frac{g_2}{2}} (e^{z_1+z_2} + e^{z_2-z_1}) + \sum_{k=2}^{n-1} g_{k+2} e^{z_{k+1}-z_k}, \quad (1.51)$$

where

$$\tilde{g}_1 = -\frac{(2ia+1)g_1}{\sqrt{2g_2}}, \quad \tilde{g}_2 = 2\left(ia + \frac{g_1}{\sqrt{2g_2}}\right) + 2\left(ia + \frac{g_1}{\sqrt{2g_2}}\right)^2. \quad (1.52)$$

Note that the coupling constants $\tilde{g}_1$ and $\tilde{g}_2$ are generic provided the parameter a is generic.

Proposition 1.4 *The integral operator with the kernel*

$$Q_{BC_n}^{I_n}(\underline{z}_n; \underline{x}_n|a) = (1 - e^{2z_{n,1}})^{-2ia} \cdot \exp \left\{ ia \sum_{i=1}^n (z_{n,i} - x_{n,i}) \right\} Q_{BC_n}^{I_n^*}(\underline{z}_n; \underline{x}_n). \quad (1.53)$$

intertwines the quadratic Hamiltonians of I_n and BC_n -type Toda chains

$$H^{I_n}(\underline{z}_n) Q_{BC_n}^{I_n}(\underline{z}_n; \underline{x}_n|a) = Q_{BC_n}^{I_n}(\underline{z}_n; \underline{x}_n|a) \cdot H^{BC_n}(\underline{x}_n). \quad (1.54)$$

Proposition 1.5 *The integral operator with the kernel*

$$Q_{BC_n}^{I_{n+1}}(\underline{z}_{n+1}; \underline{x}_n) = (1 - e^{2z_{n+1,1}})^{-2ia} \cdot \exp \left\{ ia \left(\sum_{i=1}^{n+1} z_{n+1,i} - \sum_{i=1}^n x_{n,i} \right) \right\} \cdot Q_{BC_n}^{I_{n+1}^*}(\underline{z}_n; \underline{x}_n), \quad (1.55)$$

intertwines $H^{I_{n+1}}(\underline{z}_{n+1}) - \frac{a^2}{2}$ and $H^{BC_n}(\underline{x}_n)$ Hamiltonians, that is:

$$\left(H^{I_{n+1}}(\underline{z}_{n+1}) - \frac{a^2}{2} \right) Q_{BC_n}^{I_{n+1}}(\underline{z}_{n+1}; \underline{x}_n|a) = Q_{BC_n}^{I_{n+1}}(\underline{z}_{n+1}; \underline{x}_n|a) \cdot H^{BC_n}(\underline{x}_n). \quad (1.56)$$

Let us denote the kernel of the inverse integral transformation by

$$Q_{I_k}^{BC_k}(\underline{x}_k, \underline{z}_k|a) := Q_{BC_k}^{I_k}(\underline{z}_k, \underline{x}_k|a).$$

In the following we introduce three types of iterative kernels. For any $k = 2, \dots, n$ define the integral operators $Q_{BC_k}^{BC_{k-1}}$ depending on auxiliary parameter as convolutions of the kernels of elementary intertwiners $Q_{I_k}^{BC_k}$ and $Q_{BC_{k-1}}^{I_k}$:

$$Q_{BC_{k-1}}^{BC_k}(\underline{x}_k; \underline{x}_{k-1}|\lambda_k) = \int_{\Gamma^{(k)}} Q_{I_k}^{BC_k}(\underline{x}_k; \underline{z}_k|\lambda_k) Q_{BC_{k-1}}^{I_k}(\underline{z}_k; \underline{x}_{k-1}|\lambda_k) d\underline{z}_k. \quad (1.57)$$

For $k = 1$ we set

$$Q_{BC_0}^{BC_1}(x_{1,1}|\lambda_1) = \int_{\Gamma^{(1)}} \left(1 - e^{2z_{11}} \right)^{-2i\lambda_1} e^{i\lambda_1 z_{11}} \left(\frac{1 - e^{z_{11}}}{1 + e^{z_{11}}} \right)^{\frac{2g_1}{\sqrt{2g_2}}} \exp \{ -\sqrt{g_2/2} (e^{x_{11}+z_{11}} + e^{x_{11}-z_{11}}) \} dz_{11}. \quad (1.58)$$

Let us introduce for $k = 2, \dots, n$ the integral operators $Q_{I_k}^{I_{k+1}}$

$$Q_{I_{k-1}}^{I_k}(\underline{z}_k; \underline{z}_{k-1}) = \int_{\Gamma^{(k)}} Q_{BC_{k-1}}^{I_k}(\underline{z}_k; \underline{x}_{k-1}|a) Q_{I_{k-1}}^{BC_{k-1}}(\underline{x}_{k-1}; \underline{z}_{k-1}|a) d\underline{x}_{k-1}. \quad (1.59)$$

For $k = 1$ we take

$$Q_{I_0}^{I_1}(z_{11}) = (1 + e^{z_{11}})^{-\frac{g_1}{\sqrt{2g_2}} - ia} (1 - e^{z_{11}})^{\frac{g_1}{\sqrt{2g_2}} - ia} e^{ia z_{11}}. \quad (1.60)$$

Define for $k = 2, \dots, n$ the integral operators $\Lambda_{I_k}^{I_{k+1}}$:

$$\begin{aligned} \Lambda_{I_{k-1}}^{I_k}(\underline{z}_k, \underline{z}_{k-1}|\lambda_k) = \\ \int_{\Gamma^{(1)}} \left(Q_{BC_k}^{I_k}(\underline{z}_k; \underline{x}_k|a) Q_{I_k}^{BC_k}(\underline{x}_k; \underline{u}_k|\lambda_k) \right) \left(Q_{BC_{k-1}}^{I_k}(\underline{u}_k; \underline{x}_{k-1}|\lambda_k) Q_{I_{k-1}}^{BC_{k-1}}(\underline{x}_{k-1}; \underline{u}_{k-1}|\lambda_k) \right) \\ \left(Q_{BC_{k-1}}^{I_{k-1}}(\underline{u}_{k-1}; \underline{x}_{k-1}|\lambda_k) Q_{I_{k-1}}^{BC_{k-1}}(\underline{x}_{k-1}; \underline{z}_{k-1}|a) \right) d\underline{x}_k d\underline{x}_{k-1} d\underline{u}_k d\underline{u}_{k-1}. \end{aligned} \quad (1.61)$$

For $k = 1$ we define

$$\Lambda_{I_0}^{I_1}(z_{11}|\lambda_1) = \int dx_{11} du_{11} Q_{BC_1}^{I_1}(z_{11}, x_{11}|a) Q_{I_1}^{BC_1}(x_{11}, u_{11}|\lambda_1) Q_{BC_0}^{I_1}(u_{11}|\lambda_1), \quad (1.62)$$

where

$$Q_{BC_0}^{I_1}(u_{11}|\lambda_1) = (1 + e^{u_{11}})^{-\frac{g_1}{\sqrt{2g_2}} - i\lambda_1} (1 - e^{u_{11}})^{\frac{g_1}{\sqrt{2g_2}} - i\lambda_1} e^{i\lambda_1 u_{11}}.$$

1.7 Givental representations for the eigenfunctions of BC_n and I_n Toda chains

In this Section we conjecture integral representations of elementary operators which intertwine I_n -Toda chain Hamiltonians with generic coupling constants and BC_n -Toda chain Hamiltonians with generic coupling constants. By using intertwining relations we introduce the conjectural integral formula for the eigenfunction of I_n -Toda chain with generic coupling constants corresponding to zero eigenvalues.

Conjecture 1.1 *The ground state eigenfunction for BC_n -Toda chain has the following integral representation*

$$\Psi^{BC_n}(x_1, \dots, x_n) = \int_{\Gamma} \prod_{k=1}^n Q_{BC_{k-1}}^{BC_k}(\underline{x}_k; \underline{x}_{k-1}) \prod_{k=1}^{n-1} d\underline{x}_k, \quad (1.63)$$

where $x_i := x_{n,i}$. The domain of integration Γ should be chosen in such a way that the integrand decreases exponentially at infinities.

The conjecture is also supported by various degenerations considering in the next Section. It is also easily verified for quadratic Hamiltonians using the intertwining relations (1.45), (1.47).

Conjecture 1.2 *The eigenfunction for BC_n -Toda chain corresponding to generic eigenvalues parametrized by $\underline{\lambda} = (\lambda_1, \dots, \lambda_n)$ is given by*

$$\Psi_{\underline{\lambda}}^{BC_n}(x_1, \dots, x_n) = \int_{\Gamma} \prod_{k=1}^n Q_{BC_{k-1}}^{BC_k}(\underline{x}_k; \underline{x}_{k-1}|\lambda_k) \prod_{k=1}^{n-1} d\underline{x}_k, \quad (1.64)$$

where $x_i := x_{n,i}$. The domain of integration Γ should be chosen in such a way that the integrand decreases exponentially at infinities.

It is easy to check the conjecture for quadratic Hamiltonians using the following iterative procedure:

$$H^{BC_k}(\underline{x}_k) Q_{BC_{k-1}}^{BC_k}(\underline{x}_k; \underline{x}_{k-1}|\lambda_k) = Q_{BC_{k-1}}^{BC_k}(\underline{x}_k; \underline{x}_{k-1}|\lambda_k) \left(H^{BC_{k-1}}(\underline{x}_{k-1}) + \frac{\lambda_k^2}{2} \right). \quad (1.65)$$

At the last step of the iteration we should check the relation:

$$H^{BC_1}(x_{11}) \Psi_{\lambda_1}^{BC_1}(x_{11}) = \frac{\lambda_1^2}{2} \Psi_{\lambda_1}^{BC_1}(x_{11}), \quad (1.66)$$

where

$$\Psi_{\lambda_1}^{BC_1}(x_{11}) = Q_{BC_0}^{BC_1}(x_{11}) = \int_{\Gamma(1)} \left(1 - e^{2z_{11}} \right)^{-2i\lambda_1} \left(\frac{1 - e^{z_{11}}}{1 + e^{z_{11}}} \right)^{\frac{2g_1}{\sqrt{2g_2}}} \exp\{-\sqrt{g_2/2}(e^{x_{11}+z_{11}} + e^{x_{11}-z_{11}})\} dz_{11}. \quad (1.67)$$

The last statement can be checked directly. Notice that the particular cases $g_1 = 0$ and $g_2 = 0$ corresponding to B_n and C_n -Toda chains respectively, can be transformed (up to certain multiplier depending on spectrum) to the corresponding eigenfunctions with non-zero spectrum given in [GLO2].

Conjecture 1.3 *The eigenfunction for I_n -Toda chain with generic coupling constants given by (1.51)- (1.52) corresponding to eigenvalues parametrized by $\underline{\lambda} = (\lambda_1, \dots, \lambda_n)$ has the following integral representation*

$$\Psi_{\lambda_1, \dots, \lambda_n}^{I_n}(z_1, \dots, z_n) = \int_{\Gamma} \prod_{k=1}^n \Lambda_{I_{k-1}}^{I_k}(\underline{z}_k; \underline{z}_{k-1} | \lambda_k) \prod_{k=1}^{n-1} d\underline{z}_k, \quad (1.68)$$

where $\Lambda_{I_{k-1}}^{I_k}(\underline{z}_k; \underline{z}_{k-1} | \lambda_k)$ is defined by (1.61) and $z_i := z_{n,i}$. The domain of integration Γ should be chosen in such a way that the integrand decreases exponentially at infinities.

This conjecture is easily verified for quadratic Hamiltonian using iteratively the relations

$$H^{I_k}(\underline{z}_k) \cdot \Lambda_{I_{k-1}}^{I_k}(\underline{z}_k; \underline{z}_{k-1}) = \Lambda_{I_{k-1}}^{I_k}(\underline{z}_k; \underline{z}_{k-1}) \cdot \left(H^{I_{k-1}}(\underline{z}_{k-1}) + \frac{\lambda_k^2}{2} \right), \quad (1.69)$$

and the identity

$$\left(H^{I_1}(u_{11}) - \frac{\lambda_1^2}{2} \right) Q_{BC_0}^{I_1}(u_{11} | \lambda_1) = 0.$$

1.7.1 The various limiting case of the Conjectures above are compatible with the results presented in [GLO2] and recalled in the previous Section. Let us consider a specialization $a = 0$ of the integral kernel $Q_{BC_n}^{I_n}$

$$\begin{aligned} Q_{BC_n}^{I_n^*}(\underline{z}_n; \underline{x}_n) &:= [Q_{BC_n}^{I_n}(\underline{z}_n; \underline{x}_n)]_{a=0} = \left(\frac{1 - e^{z_{n,1}}}{1 + e^{z_{n,1}}} \right)^{\frac{g_1}{\sqrt{2g_2}}} \\ &\cdot \exp \left\{ -\sqrt{\frac{g_2}{2}} \left(e^{x_{n,1}+z_{n,1}} + e^{x_{n,1}-z_{n,1}} \right) - \sum_{k=1}^{n-1} g_{k+2} e^{z_{n,k+1}-x_{n,k}} + e^{x_{n,k+1}-z_{n,k+1}} \right\}, \end{aligned} \quad (1.70)$$

that intertwines BC_n -Toda chain quadratic Hamiltonian:

$$H^{BC_n}(\underline{x}_n) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_{n,i}^2} + g_1 e^{x_{n,1}} + g_2 e^{2x_{n,1}} + \sum_{k=1}^{n-1} g_{k+2} e^{x_{n,k+1}-x_{n,k}}, \quad (1.71)$$

with the specialized I_n -Toda chain quadratic Hamiltonian:

$$\begin{aligned} H^{I_n^*}(\underline{z}_n) &= -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial z_{n,i}^2} + \frac{\tilde{g}_1}{(e^{-z_{n,1}/2} - e^{z_{n,1}/2})^2} + \frac{\tilde{g}_2}{(e^{-z_{n,1}} - e^{z_{n,1}})^2} \\ &+ g_3 \sqrt{\frac{g_2}{2}} (e^{z_{n,1}+z_{n,2}} + e^{z_{n,2}-z_{n,1}}) + \sum_{k=2}^{n-1} g_{k+2} e^{z_{n,k+1}-z_{n,k}}, \end{aligned} \quad (1.72)$$

where

$$\tilde{g}_1 = -\frac{g_1}{\sqrt{2g_2}}, \quad \tilde{g}_2 = 2 \left(\frac{g_1}{\sqrt{2g_2}} \right) \left(\frac{g_1}{\sqrt{2g_2}} + 1 \right).$$

The Hamiltonian $H^{I_n^*}(\underline{z})$ is not generic since its coupling constants $\tilde{g}_1$ and $\tilde{g}_2$ depend only on the combination $\frac{g_1}{\sqrt{2g_2}}$. Shifting the variable $z_1 \rightarrow z_1 + \frac{1}{2} \ln \frac{g_2}{2}$ one obtains the following expressions for the specialized $H^{I_n^*}(\underline{z}_n)$ and $Q_{BC_n}^{I_n^*}(\underline{z}_n; \underline{x}_n)$ when $a = 0$ (tildes over $H^{I_n^*}$ and $Q_{I_n^*}^{BC_n}$ below denote the application of the shift of variable z_1):

$$\begin{aligned} \tilde{H}^{I_n^*}(\underline{z}_n) &= -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial z_{n,i}^2} - \frac{g_1}{2} \frac{e^{-z_{n,1}} + \frac{g_2}{2} e^{z_{n,1}} - g_1}{(e^{-z_{n,1}} - \frac{g_2}{2} e^{z_{n,1}})^2} + \frac{g_2 g_3}{2} e^{z_{n,1} + z_{n,2}} + \sum_{i=1}^{n-1} g_{i+2} e^{z_{n,i+1} - z_{n,i}}, \\ \tilde{Q}_{BC_n}^{I_n^*}(\underline{z}_n, \underline{x}_n) &= \left(\frac{1 - \sqrt{\frac{g_2}{2}} e^{z_{n,1}}}{1 + \sqrt{\frac{g_2}{2}} e^{z_{n,1}}} \right)^{\frac{g_1}{\sqrt{2g_2}}} \\ &\cdot \exp \left\{ - \left(\frac{g_2}{2} e^{x_{n,1} + z_{n,1}} + e^{x_{n,1} - z_{n,1}} + \sum_{k=1}^{n-1} g_{k+2} e^{z_{n,k+1} - x_{n,k}} + e^{x_{n,k+1} - z_{n,k+1}} \right) \right\} \end{aligned}$$

Then one can directly verify the following limiting relations between various Toda chains:

1. When $g_1 \rightarrow 0$ one has

$$\begin{aligned} \lim_{g_1 \rightarrow 0} H^{BC_n}(\underline{x}_n) &= H^{C_n}(\underline{x}_n), & \lim_{g_1 \rightarrow 0} \tilde{H}^{I_n^*}(\underline{z}_n) &= H^{D_n}(\underline{z}_n), \\ \lim_{g_1 \rightarrow 0} \tilde{Q}_{BC_n}^{I_n^*}(\underline{z}_n, \underline{x}_n) &= Q_{C_n}^{D_n}(\underline{z}_n, \underline{x}_n). \end{aligned} \tag{1.73}$$

2. When $g_2 \rightarrow 0$ one obtains

$$\begin{aligned} \lim_{g_2 \rightarrow 0} H^{BC_n}(\underline{x}_n) &= H^{B_n}(\underline{x}_n), & \lim_{g_2 \rightarrow 0} \tilde{H}^{I_n^*}(\underline{z}_n) &= H^{BC_n^*}(\underline{z}_n), \\ \lim_{g_2 \rightarrow 0} \tilde{Q}_{BC_n}^{I_n^*}(\underline{z}_n, \underline{x}_n) &= Q_{B_n}^{BC_n^*}(\underline{z}_n, \underline{x}_n). \end{aligned} \tag{1.74}$$

The last formula readily follows from a simple observation:

$$\lim_{\varepsilon \rightarrow 0} \frac{g}{\sqrt{2\varepsilon}} \ln \frac{1 - \sqrt{\frac{\varepsilon}{2}} e^z}{1 + \sqrt{\frac{\varepsilon}{2}} e^z} = \lim_{\varepsilon \rightarrow 0} \frac{g}{\sqrt{2\varepsilon}} \left(-2\sqrt{\frac{\varepsilon}{2}} e^z + o(\varepsilon) \right) = -g e^z$$

2 Intertwiners for affine Toda chains

In this section we generalize the construction of the elementary intertwiners to the classical series of affine Lie algebras. For the necessary facts in the theory of affine Lie algebras see [K]. Let us first recall the construction of the Q -operator for $A_n^{(1)}$ Toda chain [PG]. Using the elementary intertwiners we propose integral operators intertwining affine Toda chains corresponding to the affine Lie algebras of the same rank.

2.1 Quadratic Hamiltonians of affine Toda chains

Let us start with a list of quadratic Hamiltonians of affine Toda chains considered below.

2.1.1 Simple roots for the system of type $A_n^{(1)}$ in the standard basis $\{e_1, \dots, e_n\}$ read as follows:

$$\alpha_0 = e_1 - e_{n+1}, \quad \alpha_i = e_{i+1} - e_i, \quad 1 \leq i \leq n.$$

The corresponding Dynkin diagram has the following form:

$$\begin{array}{c} \alpha_0 \\ \swarrow \quad \searrow \\ \alpha_1 \text{ --- } \alpha_2 \text{ --- } \dots \text{ --- } \alpha_{n-1} \text{ --- } \alpha_n \end{array} \quad (2.1)$$

Quadratic Hamiltonian operator for the $A_n^{(1)}$ closed Toda chain reads

$$\mathcal{H}^{A_n^{(1)}}(\underline{x}_{n+1}) = -\frac{1}{2} \sum_{i=1}^{n+1} \frac{\partial^2}{\partial x_{n+1,i}^2} + g_1 e^{x_{n+1,1} - x_{n+1,n+1}} + \sum_{i=1}^n g_{i+1} e^{x_{n+1,i+1} - x_{n+1,i}}. \quad (2.2)$$

2.1.2 Simple roots of the twisted affine root system $A_{2n}^{(2)}$ can be expressed in terms of the standard basis $\{e_i\}$ as follows

$$\begin{aligned} \alpha_1 &= e_1, & \alpha_{i+1} &= e_{i+1} - e_i, & 1 \leq i \leq n-1, \\ \alpha_{n+1} &= -2e_n. \end{aligned}$$

The corresponding Dynkin diagram is given by

$$\alpha_1 \Longleftarrow \alpha_2 \text{ --- } \dots \text{ --- } \alpha_{n-1} \Longleftarrow \alpha_n. \quad (2.3)$$

Introduce the affine non-reduced root system BC'_n with simple roots given by

$$\begin{aligned} \alpha_0 &= 2e_1, & \alpha_1 &= e_1, & \alpha_{i+1} &= e_{i+1} - e_i, & 1 \leq i \leq n-1, \\ \alpha_{n+1} &= -e_n - e_{n-1}, \end{aligned}$$

and the corresponding Dynkin diagram is as follows

$$\begin{array}{c} \alpha_0 \\ \parallel \\ \parallel \\ \parallel \\ \parallel \\ \parallel \\ \downarrow \\ \alpha_1 \end{array} \begin{array}{c} \searrow \\ \swarrow \end{array} \begin{array}{c} \alpha_2 \text{ --- } \dots \text{ --- } \alpha_{n-1} \end{array} \begin{array}{c} \nearrow \\ \searrow \end{array} \begin{array}{c} \alpha_n \\ \alpha_{n+1} \end{array} \quad (2.4)$$

2.1.3 Simple roots of the twisted affine root system $A_{2n-1}^{(2)}$ are given by

$$\begin{aligned} \alpha_1 &= 2e_1, & \alpha_{i+1} &= e_{i+1} - e_i, & 1 \leq i \leq n-1, \\ \alpha_{n+1} &= -e_n - e_{n-1}, \end{aligned} \quad (2.5)$$

and corresponding Dynkin diagram is

$$\begin{array}{c}
 \alpha_n \\
 \nearrow \\
 \alpha_1 \Longrightarrow \alpha_2 \text{ --- } \cdots \text{ --- } \alpha_{n-1} \\
 \searrow \\
 \alpha_{n+1}
 \end{array} \tag{2.6}$$

2.1.4 Simple roots of the affine root system $B_n^{(1)}$ are given by

$$\begin{aligned}
 \alpha_1 &= e_1, & \alpha_{i+1} &= e_{i+1} - e_i, & 1 \leq i \leq n-1, \\
 \alpha_{n+1} &= -e_n - e_{n-1},
 \end{aligned} \tag{2.7}$$

and corresponding Dynkin diagram is as follows

$$\begin{array}{c}
 \alpha_n \\
 \nearrow \\
 \alpha_1 \Longleftarrow \alpha_2 \text{ --- } \cdots \text{ --- } \alpha_{n-1} \\
 \searrow \\
 \alpha_{n+1}
 \end{array} \tag{2.8}$$

2.1.5 Introduce the affine non-reduced root system BC_n'' with simple roots given by

$$\begin{aligned}
 \alpha_0 &= 2e_1, & \alpha_1 &= e_1, & \alpha_{i+1} &= e_{i+1} - e_i, & 1 \leq i \leq n-1, \\
 \alpha_{n+1} &= -2e_n,
 \end{aligned} \tag{2.9}$$

and corresponding Dynkin diagram is given by

$$\begin{array}{c}
 \alpha_0 \\
 \parallel \\
 \parallel \\
 \parallel \\
 \parallel \\
 \parallel \\
 \parallel \\
 \downarrow \\
 \alpha_1
 \end{array}
 \begin{array}{c}
 \nearrow \\
 \searrow
 \end{array}
 \begin{array}{c}
 \alpha_2 \text{ --- } \cdots \text{ --- } \alpha_{n-1} \text{ --- } \alpha_n \Longleftarrow \alpha_{n+1}
 \end{array}$$

2.1.6 Simple roots of the affine root system $C_n^{(1)}$ are

$$\begin{aligned}
 \alpha_1 &= 2e_1, & \alpha_{i+1} &= e_{i+1} - e_i, & 1 \leq i \leq n-1, \\
 \alpha_{n+1} &= -2e_n,
 \end{aligned} \tag{2.10}$$


and corresponding Dynkin diagram is given by

$$\alpha_1 \rightrightarrows \alpha_2 \text{ --- } \dots \text{ --- } \alpha_{n-1} \text{ --- } \alpha_n \leftrightsquigarrow \alpha_{n+1} \quad .$$

2.1.7 Simple roots of the affine root system $D_n^{(1)}$ are

$$\begin{aligned} \alpha_1 &= e_1 + e_2, & \alpha_{i+1} &= e_{i+1} - e_i, & 1 \leq i \leq n-1, \\ \alpha_{n+1} &= -e_n - e_{n-1}, \end{aligned} \tag{2.11}$$

and corresponding Dynkin diagram is given by



2.1.8 Simple roots of the affine root system $D_n^{(2)}$ are

$$\begin{aligned} \alpha_1 &= e_1, & \alpha_i &= e_{i+1} - e_i, & 1 \leq i \leq n-1, \\ \alpha_{n+1} &= -e_n, \end{aligned} \tag{2.12}$$


and corresponding Dynkin diagram is given by

$$\alpha_1 \leftrightsquigarrow \alpha_1 \text{ --- } \dots \text{ --- } \alpha_n \rightrightarrows \alpha_{n+1} \quad .$$

2.1.9 Simple roots of the (non-reduced) affine root system $\widehat{BC}_{n+1}$ read as follows

$$\begin{aligned} \alpha_0 &= 2e_1 & \alpha_1 &= e_1, \\ \alpha_{i+1} &= e_{i+1} - e_i, & 1 \leq i \leq n, \\ \alpha_{n+2} &= -e_{n+1} & \alpha_{n+3} &= -2e_{n+1}, \end{aligned} \tag{2.13}$$

and corresponding Dynkin diagram is given by



Let us consider generic quadratic Toda Hamiltonian associated to the root system $\widehat{BC}_n$ (2.13):

$$\begin{aligned} \mathcal{H}^{\widehat{BC}_n}(\underline{x}_n) = & -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_{n,i}^2} + g_1 e^{x_{n,1}} + g_2 e^{2x_{n,1}} \\ & + \sum_{i=1}^{n-1} g_{i+2} e^{x_{n,i+1} - x_{n,i}} + g_{n+2} e^{-2x_{nn}} + g_{n+3} e^{-x_{nn}}. \end{aligned} \quad (2.14)$$

2.2 Elementary Intertwiners

In [PG] Pasquier-Gaudin introduce the following integral $\mathcal{Q}$ -operator defined by the integral kernel

$$Q^{A_n^{(1)}}(x_1, \dots, x_{n+1}; y_1, \dots, y_{n+1}) = \exp \left\{ - \sum_{i=1}^{n+1} \left(e^{x_i - y_i} + g_{i+1} e^{y_{i+1} - x_i} \right) \right\}, \quad (2.15)$$

where $y_{n+2} = y_1, g_{n+2} = g_1$. According to [PG] this integral kernel satisfies the following intertwining relations

$$\mathcal{H}^{A_n^{(1)}}(\underline{x}_{n+1}) \cdot Q^{A_n^{(1)}}(\underline{x}_{n+1}; \underline{y}_{n+1}) = Q^{A_n^{(1)}}(\underline{x}_{n+1}; \underline{y}_{n+1}) \cdot \mathcal{H}^{A_n^{(1)}}(\underline{y}_{n+1}).$$

Below we list kernels of the integral operators that conjecturally intertwine Hamiltonians of affine Toda chains corresponding to other classical series of (twisted) affine Lie algebras. We checked these intertwining relation against the quadratic Hamiltonians.

2.2.1 The integral operator with the following kernel

$$\begin{aligned} Q^{A_{2n}^{(2)}}_{BC'_{n+1}}(\underline{x}_n, \underline{z}_{n+1}) = & \exp \left\{ -g_1 e^{z_{n+1,1}} - \sum_{i=1}^n \left(e^{x_{n,i} - z_{n+1,i}} + g_{i+1} e^{z_{n+1,i+1} - x_{n,i}} \right) \right. \\ & \left. - g_{n+2} e^{-z_{n+1,n+1} - x_{nn}} \right\}, \end{aligned} \quad (2.16)$$

intertwines Hamiltonian operators for $A_{2n}^{(2)}$ - and $BC_{n+1}^{(2)}$ -Toda chains

$$\begin{aligned} \mathcal{H}^{A_{2n}^{(2)}}(\underline{x}_n) = & -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_{n,i}^2} + g_1 e^{x_{n,1}} + \sum_{i=1}^{n-1} g_{i+1} e^{x_{n,i+1} - x_{n,i}} + 2g_{n+1} g_{n+2} e^{-2x_{nn}}, \\ \tilde{\mathcal{H}}^{BC'_{n+1}}(\underline{z}_{n+1}) = & -\frac{1}{2} \sum_{i=1}^{n+1} \frac{\partial^2}{\partial z_{n+1,i}^2} - \frac{g_1}{2} e^{z_{n+1,1}} + \frac{g_1^2}{2} e^{2z_{n+1,1}} \\ & + \sum_{i=1}^n g_{i+1} e^{z_{n+1,i+1} - z_{n+1,i}} + g_{n+2} e^{-z_{n+1,n+1} - z_{n+1,n}}. \end{aligned} \quad (2.17)$$

The elementary intertwiner (2.16) is involutive; we use the following notation for the integral kernel of the inverse transformation:

$$Q^{BC'_{n+1}}_{A_{2n}^{(2)}}(\underline{z}_{n+1}, \underline{x}_n) := Q^{A_{2n}^{(2)}}_{BC'_{n+1}}(\underline{x}_n, \underline{z}_{n+1}). \quad (2.18)$$

2.2.2 The integral operator represented by the following kernel

$$Q_{A_{2n-1}^{(2)}}^{A_{2n-1}^{(2)}}(\underline{x}_n, \underline{z}_n) = \exp \left\{ -g_1 e^{x_{n,1}+z_{n,1}} - \sum_{i=1}^{n-1} \left(e^{x_{n,i}-z_{n,i}} + g_{i+1} e^{z_{n,i+1}-x_{n,i}} \right) - e^{x_{nn}-z_{nn}} - g_{n+1} e^{-x_{nn}-z_{nn}} \right\}, \quad (2.19)$$

intertwines Hamiltonian operators for $A_{2n-1}^{(2)}$ -Toda chains with generic coupling constants

$$\mathcal{H}^{A_{2n-1}^{(2)}}(\underline{x}_n) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_{n,i}^2} + 2g_1 e^{2x_{n,1}} + \sum_{i=1}^{n-1} g_{i+1} e^{x_{n,i+1}-x_{n,i}} + g_n g_{n+1} e^{-x_{nn}-x_{n,n-1}}, \quad (2.20)$$

$$\tilde{\mathcal{H}}^{A_{2n-1}^{(2)}}(\underline{z}_n) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial z_{n,i}^2} + g_1 g_2 e^{z_{n,1}+z_{n,2}} + \sum_{i=1}^{n-1} g_{i+1} e^{z_{n,i+1}-z_{n,i}} + 2g_{n+1} e^{-2z_{nn}}. \quad (2.21)$$

In the following we use a separate notation for integral kernel of the inverse transformation:

$$Q_{A_{2n-1}^{(2)}}^{A_{2n-1}^{(2)}}(\underline{z}_n, \underline{x}_n) := Q_{A_{2n-1}^{(2)}}^{A_{2n-1}^{(2)}}(\underline{x}_n, \underline{z}_n).$$

2.2.3 The integral operator represented by the following kernel

$$Q_{BC_n''}^{B_n^{(1)}}(\underline{x}_n, \underline{z}_n) = \exp \left\{ -g_1 e^{z_{n,1}} - \sum_{i=1}^{n-1} \left(e^{x_{n,i}-z_{n,i}} + g_{i+1} e^{z_{n,i+1}-x_{n,i}} \right) - e^{x_{nn}-z_{nn}} - g_{n+1} e^{-x_{nn}-z_{nn}} \right\}, \quad (2.22)$$

intertwines Hamiltonians of $B_n^{(1)}$ - and BC_n'' -Toda chains

$$\mathcal{H}^{B_n^{(1)}}(\underline{x}_n) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_{n,i}^2} + g_1 e^{x_{n,1}} + \sum_{i=1}^{n-1} g_{i+1} e^{x_{n,i+1}-x_{n,i}} + g_n g_{n+1} e^{-x_{nn}-x_{n,n-1}}, \quad (2.23)$$

$$\tilde{\mathcal{H}}^{BC_n''}(\underline{z}_n) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial z_{n,i}^2} - \frac{g_1}{2} e^{z_{n,1}} + \frac{g_1^2}{2} e^{2z_{n,1}} + \sum_{i=1}^{n-1} g_{i+1} e^{z_{n,i+1}-z_{n,i}} + 2g_{n+1} e^{-2z_{nn}}. \quad (2.24)$$

We use the following notation for integral kernel of the inverse transformation:

$$Q_{B_n^{(1)}}^{BC_n''}(\underline{z}_n, \underline{x}_n) := Q_{BC_n''}^{B_n^{(1)}}(\underline{x}_n, \underline{z}_n). \quad (2.25)$$

2.2.4 The integral operator with the following kernel

$$Q_{D_{n+1}}^{C_n^{(1)}}(\underline{x}_n, \underline{z}_{n+1}) = \exp \left\{ -g_1 e^{x_{n,1}+z_{n+1,1}} - \sum_{i=1}^n \left(e^{x_{n,i}-z_{n+1,i}} + g_{i+1} e^{z_{n+1,i+1}-x_{n,i}} \right) - g_{n+2} e^{-z_{n+1,n+1}-x_{nn}} \right\} \quad (2.26)$$

intertwines Hamiltonian operators for $C_n^{(1)}$ - and $D_{n+1}^{(1)}$ -Toda chains

$$\mathcal{H}^{C_n^{(1)}}(\underline{x}_n) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_{n,i}^2} + 2g_1 e^{2x_{n,1}} + \sum_{i=1}^{n-1} g_{i+1} e^{x_{n,i+1}-x_{n,i}} + 2g_{n+1} g_{n+2} e^{-2x_{nn}}, \quad (2.27)$$

$$\mathcal{H}^{D_{n+1}^{(1)}}(\underline{z}_{n+1}) = -\frac{1}{2} \sum_{i=1}^{n+1} \frac{\partial^2}{\partial z_{n+1,i}^2} + g_1 g_2 e^{z_{n+1,1}+z_{n+1,2}} + \sum_{i=1}^n g_{i+1} e^{z_{n+1,i+1}-z_{n+1,i}} + g_{n+2} e^{-z_{n+1,n+1}-z_{n+1,n}}. \quad (2.28)$$

2.2.5 The integral operator with the kernel

$$Q_{C_{n-1}}^{D_n^{(1)}}(\underline{x}_n, \underline{z}_{n-1}) = \exp \left\{ -g_1 e^{x_{n,1}+z_{n-1,1}} - \sum_{i=1}^{n-1} \left(e^{z_{n-1,i}-x_{n,i}} + g_{i+1} e^{x_{n,i+1}-z_{n-1,i}} \right) - g_{n+1} e^{-x_{nn}-z_{n-1,n-1}} \right\}, \quad (2.29)$$

intertwines Hamiltonian operators for $D_n^{(1)}$ - and $C_{n-1}^{(1)}$ -Toda chains

$$\mathcal{H}^{D_n^{(1)}}(\underline{x}_n) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_{n,i}^2} + g_1 g_2 e^{x_{n,1}+x_{n,2}} + \sum_{i=1}^{n-1} g_{i+1} e^{x_{n,i+1}-x_{n,i}} + g_{n+1} e^{-x_{nn}-x_{n,n-1}}, \quad (2.30)$$

$$\mathcal{H}^{C_{n-1}^{(1)}}(\underline{z}_{n-1}) = -\frac{1}{2} \sum_{i=1}^{n-1} \frac{\partial^2}{\partial z_{n-1,i}^2} + 2g_1 e^{2z_{n-1,1}} + \sum_{i=1}^{n-2} g_{i+1} e^{z_{n-1,i+1}-z_{n-1,i}} + 2g_n g_{n+1} e^{-2z_{n-1,n-1}}. \quad (2.31)$$

We use the following notations for integral kernels of the inverse transformations:

$$Q_{C_n^{(1)}}^{D_{n+1}^{(1)}}(\underline{z}_{n+1}, \underline{x}_n) := Q_{D_{n+1}}^{C_n^{(1)}}(\underline{x}_n, \underline{z}_{n+1}), \quad (2.32)$$

$$Q_{D_n^{(1)}}^{C_{n-1}^{(1)}}(\underline{z}_{n-1}, \underline{x}_n) := Q_{C_{n-1}}^{D_n^{(1)}}(\underline{x}_n, \underline{z}_{n-1}).$$

2.2.6 The integral operator with the following kernel

$$Q_{\widehat{BC}_{n+1}}^{D_n^{(2)}}(\underline{x}_n, \underline{z}_{n+1}) = \exp \left\{ -g_1 e^{z_{n+1,1}} - \sum_{i=1}^n \left(e^{x_{n,i}-z_{n+1,i}} + g_{i+1} e^{z_{n+1,i+1}-x_{n,i}} \right) - g_{n+2} e^{-z_{n+1,n+1}} \right\} \quad (2.33)$$

intertwines Hamiltonian operators for $D_n^{(2)}$ - and $\widehat{BC}_{n+1}$ -Toda chains defined as:

$$\mathcal{H}^{D_n^{(2)}}(\underline{x}_n) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_{n,i}^2} + g_1 e^{x_1} + \sum_{i=1}^{n-1} g_{i+1} e^{x_{n,i+1} - x_{n,i}} + g_{n+1} e^{-x_{nn}}, \quad (2.34)$$

$$\begin{aligned} \mathcal{H}^{\widehat{BC}_{n+1}^*}(\underline{z}_{n+1}) &= -\frac{1}{2} \sum_{i=1}^{n+1} \frac{\partial^2}{\partial z_{n+1,i}^2} - \frac{g_1}{2} e^{z_{n+1,1}} + \frac{g_1^2}{2} e^{2z_{n+1,1}} \\ &+ \sum_{i=1}^n g_{i+1} e^{z_{n+1,i+1} - z_{n+1,i}} - \frac{g_{n+2}}{2} e^{-z_{n+1,n+1}} + \frac{g_{n+2}^2}{2} e^{-2z_{n+1,n+1}}. \end{aligned} \quad (2.35)$$

The integral kernel for the inverse transformation is denoted by

$$Q_{D_n^{(2)}}^{\widehat{BC}_{n+1}^*}(\underline{z}_{n+1}, \underline{x}_n) := Q_{\widehat{BC}_{n+1}^*}^{D_n^{(2)}}(\underline{x}_n, \underline{z}_{n+1}). \quad (2.36)$$

Proposition 2.1 *Integral operator with the kernel*

$$\begin{aligned} Q_{\widehat{I}_{n+1}}^{\widehat{BC}_n}(\underline{z}_{n+1}; \underline{x}_n) &= \left(\frac{1 - e^{z_{n+1,1}}}{1 + e^{z_{n+1,1}}} \right)^{\frac{g_1}{\sqrt{2}g_2}} \left(\frac{1 - e^{z_{n+1,n+1}}}{1 + e^{z_{n+1,n+1}}} \right)^{\frac{g_{n+3}}{\sqrt{2}g_{n+2}}} \left(\frac{1 - e^{-2z_{n+1,n+1}}}{1 - e^{2z_{n+1,1}}} \right)^a \\ &\cdot \exp \left\{ a \left(\sum_{i=1}^{n+1} z_{n+1,i} - \sum_{k=1}^n x_{n,k} \right) - \sqrt{\frac{g_2}{2}} \left(e^{x_{n,1} + z_{n+1,1}} + e^{x_{n,1} - z_{n+1,1}} \right) \right. \\ &\quad - \sum_{k=1}^{n-1} g_{k+2} e^{z_{n+1,k+1} - x_{n,k}} + e^{x_{n,k+1} - z_{n,k}} \\ &\quad \left. - \sqrt{\frac{g_{n+2}}{2}} \left(e^{z_{n+1,n+1} - x_{nn}} + e^{-z_{n+1,n+1} - x_{nn}} \right) \right\}, \end{aligned} \quad (2.37)$$

intertwines quadratic Hamiltonians of the BC_n -Toda chain:

$$\begin{aligned} H^{\widehat{BC}_n}(\underline{x}_n) &= -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_{n,i}^2} + g_1 e^{x_{n,1}} + g_2 e^{2x_{n,1}} + \sum_{k=1}^{n-1} g_{k+2} e^{x_{n,k+1} - x_{n,k}} \\ &\quad + g_{n+2} e^{-2x_{nn}} - g_{n+3} e^{-x_{nn}}, \end{aligned}$$

with the closed Inozemtsev quadratic Hamiltonian:

$$\begin{aligned} H^{\widehat{I}_{n+1}}(\underline{z}_{n+1}) &= -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial z_{n+1,i}^2} + \frac{\tilde{g}_1}{(e^{-z_{n+1,1}/2} - e^{z_{n+1,1}/2})^2} + \frac{\tilde{g}_2}{(e^{-z_{n+1,1}} - e^{z_{n+1,1}})^2} \\ &+ g_3 \sqrt{\frac{g_2}{2}} (e^{z_{n+1,1} + z_{n+1,2}} + e^{z_{n+1,2} - z_{n+1,1}}) + \sum_{k=2}^{n-1} g_{k+2} e^{z_{n+1,k+1} - z_{n+1,k}} \\ &\quad + \sqrt{\frac{g_{n+2}}{2}} (e^{z_{n+1,1} + z_{n+1,2}} + e^{z_{n+1,2} - z_{n+1,1}}) + \frac{a^2}{2} \\ &\quad + \frac{\tilde{g}_{n+3}}{(e^{-z_{n+1,n+1}} - e^{z_{n+1,n+1}})^2} + \frac{\tilde{g}_{n+4}}{(e^{-z_{n+1,n+1}/2} - e^{z_{n+1,n+1}/2})^2}, \end{aligned} \quad (2.38)$$

where

$$\begin{aligned}\tilde{g}_1 &= -\frac{(2a+1)g_1}{\sqrt{2g_2}}, & \tilde{g}_2 &= 2\left(a + \frac{g_1}{\sqrt{2g_2}}\right) + 2\left(a + \frac{g_1}{\sqrt{2g_2}}\right)^2, \\ \tilde{g}_{n+3} &= 2\left(a + \frac{g_{n+3}}{\sqrt{2g_{n+2}}}\right)^2 - 2\left(a - \frac{g_{n+3}}{\sqrt{2g_{n+2}}}\right), & \tilde{g}_{n+4} &= (2a-1)\frac{g_{n+3}}{\sqrt{2g_{n+2}}}.\end{aligned}\tag{2.39}$$

2.3 Q -operators for affine Toda chains

Now we apply the results presented in the previous Sections to the construction of the integral Q -operators for all classical series of affine Lie algebras as well as affine versions of BC_n - and I_n -Toda chains. Let us note that the elementary intertwining operators and recursive operators for finite Lie algebras can be obtained from the elementary intertwining operators and Q -operators for affine Lie algebras by taking appropriate limits $g_i \rightarrow 0$ in (2.16)-(2.29). This generalizes known relation between Q -operator for $A_n^{(1)}$ and recursive operators for A_n .

Conjecture 2.1 *The integral kernels for Q -operators have the following form*

$$Q^{A_{2n}^{(2)}}(\underline{x}; \underline{y}) = \int Q^{A_{2n}^{(2)}}_{BC'_{n+1}}(\underline{x}; \underline{z}_{n+1}) Q^{BC'_{n+1}}_{A_{2n}^{(2)}}(\underline{z}_{n+1}; \underline{y}) d\underline{z}_{n+1}, \tag{2.40}$$

$$Q^{A_{2n-1}^{(2)}}(\underline{x}; \underline{y}) = \int Q^{A_{2n-1}^{(2)}}_{A_{2n-1}^{(2)}}(\underline{x}; \underline{z}_n) Q^{A_{2n-1}^{(2)}}_{A_{2n-1}^{(2)}}(\underline{z}_n; \underline{y}) d\underline{z}_n, \tag{2.41}$$

$$Q^{B_n^{(1)}}(\underline{x}; \underline{y}) = \int Q^{B_n^{(1)}}_{BC''_n}(\underline{x}; \underline{z}_n) Q^{BC''_n}_{B_n^{(1)}}(\underline{z}_n; \underline{y}) d\underline{z}_n, \tag{2.42}$$

$$Q^{C_n^{(1)}}(\underline{x}; \underline{y}) = \int Q^{C_n^{(1)}}_{D_{n+1}^{(1)}}(\underline{x}; \underline{z}_{n+1}) Q^{D_{n+1}^{(1)}}_{C_n^{(1)}}(\underline{z}_{n+1}; \underline{y}) d\underline{z}_{n+1}, \tag{2.43}$$

$$Q^{D_n^{(1)}}(\underline{x}; \underline{y}) = \int Q^{D_n^{(1)}}_{C_{n-1}^{(1)}}(\underline{x}; \underline{z}_{n-1}) Q^{C_{n-1}^{(1)}}_{D_n^{(1)}}(\underline{z}_{n-1}; \underline{y}) d\underline{z}_{n-1}, \tag{2.44}$$

$$Q^{D_n^{(2)}}(\underline{x}; \underline{y}) = \int Q^{D_n^{(2)}}_{\widehat{BC}_{n+1}}(\underline{x}; \underline{z}_{n+1}) Q^{\widehat{BC}_{n+1}}_{D_n^{(2)}}(\underline{z}_{n+1}; \underline{y}) d\underline{z}_{n+1}. \tag{2.45}$$

$$Q^{\widehat{BC}_n}(\underline{x}; \underline{y}) = \int Q^{\widehat{BC}_n}_{\widehat{I}_{n+1}}(\underline{x}; \underline{z}_{n+1}) Q^{\widehat{I}_{n+1}}_{\widehat{BC}_n}(\underline{z}_{n+1}; \underline{y}) d\underline{z}_{n+1}, \tag{2.46}$$

$$Q^{\widehat{I}_n}(\underline{x}; \underline{y}) = \int Q^{\widehat{I}_n}_{\widehat{BC}_{n-1}}(\underline{x}; \underline{z}_{n-1}) Q^{\widehat{BC}_{n-1}}_{\widehat{I}_n}(\underline{z}_{n-1}; \underline{y}) d\underline{z}_{n-1}, \tag{2.47}$$

where $\underline{x} = (x_1, \dots, x_n)$, $\underline{y} = (y_1, \dots, y_n)$, and $\underline{z}_k = (z_{k,1}, \dots, z_{k,k})$ for $k = n, n \pm 1$.

3 Intertwining operators semi-infinite Toda chains

Similar approach can be applied to construct Q -operators for infinite root systems B_∞ , C_∞ , BC_∞ and D_∞ .

3.1 Quadratic Hamiltonians for semi-infinite Toda chains

Simple roots and Dynkin diagrams for infinite Lie algebras A_∞ , B_∞ , C_∞ , BC_∞ and D_∞ are as follows

$$A_\infty : \quad \alpha_{i+1} = e_{i+1} - e_i, \quad i \in \mathbb{Z}, \quad (3.1)$$

$$\dots \text{ --- } \alpha_{-1} \text{ --- } \alpha_0 \text{ --- } \alpha_1 \text{ --- } \alpha_2 \text{ --- } \dots$$

$$B_\infty : \quad \alpha_1 = e_1, \quad \alpha_{i+1} = e_{i+1} - e_i, \quad i \in \mathbb{Z}_{>0}, \quad (3.2)$$

$$\alpha_1 \Longleftarrow \alpha_2 \text{ --- } \alpha_3 \text{ --- } \dots$$

$$C_\infty : \quad \alpha_1 = 2e_1, \quad \alpha_{i+1} = e_{i+1} - e_i, \quad i \in \mathbb{Z}_{>0}, \quad (3.3)$$

$$\alpha_1 \Longrightarrow \alpha_2 \text{ --- } \alpha_3 \text{ --- } \dots$$

$$D_\infty : \quad \alpha_1 = e_1 + e_2, \quad \alpha_{i+1} = e_{i+1} - e_i, \quad i \in \mathbb{Z}_{>0}, \quad (3.4)$$

$$\begin{array}{c} \alpha_1 \\ \diagdown \\ \alpha_3 \text{ --- } \alpha_4 \text{ --- } \dots \\ \diagup \\ \alpha_2 \end{array}$$

$$BC_\infty : \quad \alpha_0 = 2e_1, \quad \alpha_1 = e_1, \quad \alpha_{i+1} = e_{i+1} - e_i, \quad i \in \mathbb{Z}_{>0}, \quad (3.5)$$

$$\begin{array}{c} \alpha_0 \\ \parallel \\ \parallel \\ \parallel \\ \parallel \\ \parallel \\ \parallel \\ \downarrow \\ \alpha_1 \end{array} \begin{array}{c} \searrow \\ \searrow \end{array} \begin{array}{c} \alpha_2 \text{ --- } \alpha_3 \text{ --- } \dots \end{array}$$

3.2 Elementary intertwiners for semi-infinite Toda chains

Q -operator for A_∞ infinite Toda chain is known (see [T] for the classical limit) and has an elementary structure. Namely, it is given by an integral operator with the kernel

$$Q(\underline{x}, \underline{y}) = \exp \left\{ - \sum_{i \in \mathbb{Z}} \left(e^{x_i - y_i} + g_i e^{y_{i+1} - x_i} \right) \right\}, \quad (3.6)$$

and intertwines A_∞ Toda chain Hamiltonians e.g.

$$\mathcal{H}^{A_\infty}(\underline{x}) = -\frac{1}{2} \sum_{i \in \mathbb{Z}} \frac{\partial^2}{\partial x_i^2} + \sum_{i \in \mathbb{Z}} g_i e^{x_{i+1} - x_i}, \quad (3.7)$$

$$\mathcal{H}^{A_\infty}(\underline{y}) = -\frac{1}{2} \sum_{i \in \mathbb{Z}} \frac{\partial^2}{\partial y_i^2} + \sum_{i \in \mathbb{Z}} g_i e^{y_{i+1} - y_i}. \quad (3.8)$$

In this section the notations $\underline{x} = (x_1, x_2, \dots)$, $\underline{y} = (y_1, y_2, \dots)$ are used. Below we conjecture a generalization of this integral operator to other classical series as well as BC and I series of Toda chains. The intertwining relations with quadratic Hamiltonians can be checked straightforwardly.

3.2.1 The integral operator with the kernel

$$Q_{BC_\infty}^{B_\infty}(\underline{x}, \underline{z}) = \exp \left\{ - g_1 e^{z_1} - \sum_{i > 0} \left(e^{x_i - z_i} + g_{i+1} e^{z_{i+1} - x_i} \right) \right\}, \quad (3.9)$$

intertwines B_∞ and BC_∞^* Toda chain Hamiltonian operators

$$\mathcal{H}^{B_\infty}(\underline{x}) = -\frac{1}{2} \sum_{i=1}^{\infty} \frac{\partial^2}{\partial x_i^2} + g_1 e^{x_1} + \sum_{i=1}^{\infty} g_{i+1} e^{x_{i+1} - x_i}, \quad (3.10)$$

$$\mathcal{H}^{BC_\infty^*}(\underline{z}) = -\frac{1}{2} \sum_{i=1}^{\infty} \frac{\partial^2}{\partial z_i^2} - \frac{g_1}{2} e^{z_1} + \frac{g_1^2}{2} e^{2z_1} + \sum_{i=1}^{\infty} g_{i+1} e^{z_{i+1} - z_i}. \quad (3.11)$$

3.2.2 Similarly the integral operator with the kernel

$$Q_{D_\infty}^{C_\infty}(\underline{x}, \underline{z}) = \exp \left\{ - g_1 e^{x_1 + z_1} - \sum_{i > 0} \left(e^{x_i - z_i} + g_{i+1} e^{z_{i+1} - x_i} \right) \right\}, \quad (3.12)$$

intertwines C_∞ and D_∞ Toda chain Hamiltonian operators

$$\mathcal{H}^{C_\infty}(\underline{x}) = -\frac{1}{2} \sum_{i=1}^{\infty} \frac{\partial^2}{\partial x_i^2} + 2g_1 e^{2x_1} + \sum_{i=1}^{\infty} g_{i+1} e^{x_{i+1} - x_i}, \quad (3.13)$$

$$\mathcal{H}^{D_\infty}(\underline{z}) = -\frac{1}{2} \sum_{i=1}^{\infty} \frac{\partial^2}{\partial z_i^2} + g_1 g_2 e^{z_1 + z_2} + \sum_{i=1}^{\infty} g_{i+1} e^{z_{i+1} - z_i}. \quad (3.14)$$

Thus integral Q -operator for C_∞ has as the kernel

$$Q^{C_\infty}(\underline{x}; \underline{y}) = \int Q_{D_\infty}^{C_\infty}(\underline{x}, \underline{z}) \cdot Q_{C_\infty}^{D_\infty}(\underline{z}, \underline{y}) \prod_{i=1}^{\infty} dz_i.$$

3.2.3 The integral operator with the kernel

$$Q_{C_\infty}^{D_\infty}(\underline{z}, \underline{x}) = \exp \left\{ -g_1 e^{x_1+z_1} - \sum_{i>0} \left(e^{z_i-x_i} + g_{i+1} e^{x_{i+1}-z_i} \right) \right\}, \quad (3.15)$$

intertwines D_∞ and C_∞ Toda chain Hamiltonian operators

$$\mathcal{H}^{D_\infty}(\underline{x}) = -\frac{1}{2} \sum_{i=1}^{\infty} \frac{\partial^2}{\partial x_i^2} + g_1 g_2 e^{x_1+x_2} + \sum_{i=1}^{\infty} g_{i+1} e^{x_{i+1}-x_i}, \quad (3.16)$$

$$\mathcal{H}^{C_\infty}(z_i) = -\frac{1}{2} \sum_{i=1}^{\infty} \frac{\partial^2}{\partial z_i^2} + 2g_1 e^{2z_1} + \sum_{i=1}^{\infty} g_{i+1} e^{z_{i+1}-z_i}. \quad (3.17)$$

3.2.4 Let us consider the quadratic Hamiltonian of BC_∞ -Toda chain with arbitrary coupling constants:

$$\mathcal{H}^{BC_\infty}(\underline{x}) = -\frac{1}{2} \sum_{i=1}^{\infty} \frac{\partial^2}{\partial x_i^2} + g_1 e^{x_1} + g_2 e^{2x_1} + \sum_{i=1}^{\infty} g_{i+2} e^{x_{i+1}-x_i}.$$

Then an integral operator with the kernel

$$\begin{aligned} Q_{I_\infty}^{BC_\infty}(\underline{x}; \underline{z}) &= (1 + e^{z_{n,1}})^{-\frac{g_1}{\sqrt{2g_2}}-a} (1 - e^{z_{n,1}})^{\frac{g_1}{\sqrt{2g_2}}-a} \cdot \exp \left\{ a \sum_{i>0} (z_{n,i} - x_{n,i}) \right. \\ &\quad \left. - \sqrt{\frac{g_2}{2}} \left(e^{x_1+z_1} + e^{x_1-z_1} \right) - \sum_{i>0} \left(g_{i+1} e^{z_{i+1}-x_i} + e^{x_{i+1}-z_{i+1}} \right) \right\} \end{aligned} \quad (3.18)$$

intertwines $\mathcal{H}^{BC_\infty}(\underline{x})$ with the specified quadratic Hamiltonian of I_∞ integrable system:

$$\begin{aligned} H^{I_\infty}(\underline{z}) &= -\frac{1}{2} \sum_{i=1}^{\infty} \frac{\partial^2}{\partial z_i^2} + \frac{\tilde{g}_1}{(e^{-z_1/2} - e^{z_1/2})^2} + \frac{\tilde{g}_2}{(e^{-z_1} - e^{z_1})^2} \\ &\quad + \sqrt{\frac{g_2}{2}} e^{z_1+z_2} + g_3 \sqrt{\frac{g_2}{2}} e^{z_2-z_1} + \sum_{i=2}^{\infty} g_{i+2} e^{z_{i+1}-z_i}, \end{aligned} \quad (3.19)$$

where the coupling constants $\tilde{g}_1$ and $\tilde{g}_2$ are given by (1.52):

$$\tilde{g}_1 = -\frac{(2a+1)g_1}{\sqrt{2g_2}}, \quad \tilde{g}_2 = 2\left(a + \frac{g_1}{\sqrt{2g_2}}\right) + 2\left(a + \frac{g_1}{\sqrt{2g_2}}\right)^2.$$

Let us denote by $Q_{BC_\infty}^{I_\infty}(\underline{z}; \underline{x})$ the kernel of the inverse integral transformation.

3.3 Q -operators for semi-infinite Toda chains

Now one can construct Q -operators for the semi-infinite Toda chains, combining the elementary intertwiners obtained in previous Subsections.

Conjecture 3.1 *The Q -operators for semi-infinite Toda chains are given by the following integral kernels.*

$$Q^{B_\infty}(\underline{x}; \underline{y}) = \int Q_{BC_\infty}^{B_\infty}(x_i, z_i) \cdot Q_{BC_\infty}^{BC_\infty}(z_i, y_i) \prod_{i=1}^{\infty} dz_i; \quad (3.20)$$

$$Q^{C\infty}(\underline{x}; \underline{y}) = \int Q^{C\infty}_{D\infty}(\underline{x}, \underline{z}) \cdot Q^{D\infty}_{C\infty}(\underline{z}, \underline{y}) \prod_{i=1}^{\infty} dz_i; \quad (3.21)$$

$$Q^{D\infty}(\underline{x}; \underline{y}) = \int Q^{D\infty}_{C\infty}(x_i, z_i) \cdot Q^{C\infty}_{D\infty}(z_i, y_i) \prod_{i=1}^{\infty} dz_i; \quad (3.22)$$

$$Q^{BC\infty}(\underline{x}; \underline{y}) = \int Q^{BC\infty}_{I\infty}(x_i, z_i) \cdot Q^{I\infty}_{BC\infty}(z_i, y_i) \prod_{i=1}^{\infty} dz_i; \quad (3.23)$$

$$Q^{I\infty}(\underline{z}; \underline{y}) = \int Q^{I\infty}_{BC\infty}(z_i, x_i) \cdot Q^{BC\infty}_{I\infty}(x_i, y_i) \prod_{i=1}^{\infty} dx_i. \quad (3.24)$$

One can directly check this conjecture for the intertwining relations with quadratic Hamiltonians.

References

- [F] L.D. Faddeev, *How algebraic Bethe ansatz works for integrable models*, Proceedings of Les Houches School of Physics: Quantum Symmetries, Les Houches, (1995), 149–219.
- [GKLO] A. Gerasimov, S. Kharchev, D. Lebedev, S. Oblezin, *On a Gauss-Givental representation for quantum Toda chain wave function*, Int. Math. Res. Notices, Volume 2006 (2006), ArticleID 96489, [arXiv:math.RT/0505310](#).
- [GLO1] A. Gerasimov, D. Lebedev, S. Oblezin, *Givental Integral Representation for Classical Groups*, [arXiv:math. RT/ 0608152](#).
- [GLO2] A. Gerasimov, D. Lebedev, S. Oblezin, *New integral representations of Whittaker functions for classical groups*, [math.RT/0705.2886](#).
- [GLO3] A. Gerasimov, D. Lebedev, S. Oblezin, *Baxter Operator and Archimedean Hecke Algebra*, Commun. Math. Physics. DOI 10.1007/s00220-008-0547-9, [arXiv:math.RT/0706.3476](#).
- [Gi] A. Givental, *Stationary Phase Integrals, Quantum Toda Lattices, Flag Manifolds and the Mirror Conjecture*, AMS Trans. (2) **180** (1997), 103–115, [arXiv:alg-geom/9612001](#).
- [Ha] M. Hashizume, *Whittaker functions on semi-simple Lie groups*, Hiroshima Math.J., **12**, (1982), 259–293.
- [He] S. Helgason, *Differential geometry, Lie groups, and symmetric spaces*, **v.34**, Graduated studies in Mathematics, American Mathematical Society, 1978.
- [I] V. Inozemtsev, *The finite Toda lattices*, Commun. Math. Phys. **121** (1989), 629–638.
- [K] V. Kac, *Infinite-dimensional Lie algebras*, Cambridge University Press, Cambridge 1990.
- [KS] V. Kuznetsov, E.K. Sklyanin, *Backlund Transformation for the BC-Type Toda Lattice*, Vadim Kuznetsov Memorial Issue on Integrable Systems and Related Topics, Symmetry, Integrability and Geometry: Methods and Applications, SIGMA 3 (2007), 080, 17 pages; [arXiv:0707.1950](#).

- [PG] V. Pasquier, M. Gaudin, *The periodic Toda chain and a matrix generalization of the Bessel function recursion relation*, J. Phys. A **25** (1992), 5243–5252.
- [RSTS] A.G. Reyman, M.A. Semenov-Tian-Shansky, *Integrable Systems. Group theory approach*, Modern Mathematics, Moscow-Igevs: Institute Computer Sciences, 2003.
- [S] E.K. Sklyanin, *Boundary conditions for integrable quantum systems*, J. Phys. A: Math. Gen. **21** (1988), 2375–389.
- [STS] M.A. Semenov-Tian-Shansky, *Quantization of Open Toda Lattices*, Encyclopædia of Mathematical Sciences, vol. 16. Dynamical Systems VII. Ch. 3. Springer Verlag, 1994, 226–259.
- [T] M. Toda, *Theory of Nonlinear Lattices*, Berlin, Springer-Verlag, 1981.

A.G. *Institute for Theoretical and Experimental Physics, 117259, Moscow, Russia;*
School of Mathematics, Trinity College, Dublin 2, Ireland;
Hamilton Mathematics Institute, Trinity College, Dublin 2, Ireland;

D.L. *Institute for Theoretical and Experimental Physics, 117259, Moscow, Russia;*
E-mail address: lebedev@itep.ru

S.O. *Institute for Theoretical and Experimental Physics, 117259, Moscow, Russia;*
E-mail address: Sergey.Oblezin@itep.ru